



LIGHT W/O EMULSION 6346F

Formula

A	• MONTANE 481VG (<i>Sorbitan oleate/Beeswax/Stearic acid - SEPPIC</i>)	5.00 %
	• PEG-45 Dodecylglycol copolymer	1.00 %
	• Cetearyl octanoate	14.00 %
	• Paraffin oil	8.00 %
	• SEPICIDE HB (<i>Phenoxyethanol/Methylparaben/Ethylparaben/Propylparaben/Butyl paraben - SEPPIC</i>)	1.00 %
B	• Water	QSP 100 %
	• Glycerin	5.00 %
	• MgSO4 7H2O	0.70 %
	• MICROPEARL MHB (<i>Polymethylmetacrylate - SEPPIC</i>)	1.00 %
C	• Fragrance	0.20 %

Procedure

Heat A and B to 80°C separately . The MICROPEARL must be added to the heated aqueous phase just before emulsification. Then emulsify **B into A** using a sufficient rate of shear; Maintain the shear rate until the emulsion has cooled down then add perfume at 30°C.

Comments

MONTANE 481VG A water in oil emulsifier which allows emulsification of low polar oils like paraffin oil or squalane. It provides the emulsions with a great stability: no oily exsudation after 3 months at 50°C. The emulsion is also resistant to freeze-thaw cycles (-5°C/+40°C)

MICROPEARL M100 A fine powder with an outstanding velvety feel. It significantly reduces the typical after-feel of water-in-oil emulsions.

SEPICIDE HB Preservative

Characteristics

appearance Smooth white cream
viscosity About 50,000 mPa.s BROOKFIELD LVT S4 6rpm
stability Excellent at room temperature /40°C/50°C
Stable after freeze-thaw cycles (-5°C/+40°C)
Stable after centrifugation at 50°C



Notes

PEG-45 Dodecylglycol copolymer: ELFACOS ST9 (AKZO)
Fragrance: NIVE G92.27190 (ROBERTET)

6346F - SEPPIC - A9602

Since the proposed formulation has not undergone a toxicological study, the handling and use of the proposed products are given as an indication only and in no way bind SEPPIC's responsibility.